

PROPER USE OF THIS MEDICATION

Pr Taro-Clobetasol Spray

Clobetasol propionate solution, 0.05%

Usual dose:

Apply twice daily, once in the morning and once at night. Use only enough to cover the affected areas. Make sure that you use Taro-Clobetasol Spray exactly as directed by your doctor.

Before applying Taro-Clobetasol Spray, wash the area to be treated with a mild cleanser, pat dry, and wait several minutes.

How to use Taro-Clobetasol Spray:

Please read the following instructions before using Taro-Clobetasol (clobetasol propionate) Spray, 0.05%. The terms described in the figure below (Fig.1) will help you understand these instructions.

When you receive Taro-Clobetasol Spray the Directional Spray Nozzle is in the “locked” position (see Fig. 2).

Fig. 1




Fig.2



To use Taro-Clobetasol Spray follow Steps 1 through 3.

Step 1: Grip the sides of the Pump Top with one hand and use your second hand to point the Directional Spray Nozzle where you want the spray to go (see Fig. 3). The spray will be delivered through the nozzle opening at the end of the Directional Spray Nozzle.

Fig. 3




Step 2: Push down on the Pump Top to spray Taro-Clobetasol Spray (see Fig.4).

Fig. 4



Step 3: Spray only enough to cover affected area. Rub gently to ensure even coverage. Do not apply Taro-Clobetasol Spray to your face, underarms or groin and avoid contact with eyes and lips (see Fig. 5).

Fig. 5



The maximum recommended single dose should not exceed 3.6 mL (about $\frac{3}{4}$ of a teaspoon), or 23 sprays (each pump spray is about 0.16 mL). Single application not to exceed 20% body surface area.

Wash your hands after applying Taro-Clobetasol Spray.

DO NOT APPLY MORE THAN THE PRESCRIBED AMOUNT (50 g or one complete bottle of spray) per week maximum.

Not to be used with occlusive dressings.

Stop using Taro-Clobetasol Spray as soon as your plaques have healed. After four weeks, you must stop using the product, even if you are not completely healed, and speak to your doctor.

Taro-Clobetasol Spray is for external use only. Avoid getting it in your eyes and mouth.

Taro-Clobetasol Spray is flammable. Avoid using it near sources of heat or open flame.

Discard product 28 days after receipt from pharmacist.

This leaflet was prepared by Taro Pharmaceuticals Inc.
Taro-Clobetasol Spray Product Monograph
Last Prepared: June 14, 2019



COMMITTED TO GROWTH. DEDICATED TO QUALITY.

1.800.268.1975 | www.taro.ca